

# Y-Tuss

Chlophedianol Hydrochloride 12.5 mg

Brompheniramine Maleate 2 mg

Pseudoephedrine Hydrochloride 30 mg



**Temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:**

- ✓ cough due to minor throat and bronchial irritation
- ✓ nasal congestion
- ✓ reduces swelling of nasal passages
- ✓ runny nose
- ✓ sneezing
- ✓ itching of the nose or throat
- ✓ itchy, watery eyes

- Unique combination of non-narcotic cough suppressant, antihistamine, and decongestant
- Centrally and locally acting cough suppressant that relieves irritated throat
- Alcohol Free, Dye Free, Sugar Free, and Gluten Free
- Approximate retail price range is \$16-\$19 per 4 oz.

## DOSAGE - TID

- **Adults and Children over 12**  
2 tsp. every 8 hours
- **Children 6-12**  
1 tsp. every 8 hours

***Delicious  
Grape Flavor***

**MAGNA**  
Pharmaceuticals, Inc.  
*Accountability*

# Y-Tuss

NDC 58407-302-04

## Drug Facts

**Active ingredients** (in each 5 mL teaspoonful)

**Brompheniramine Maleate** 2 mg - Antihistamine

**Chlophedianol Hydrochloride** 12.5 mg - Cough Suppressant

**Pseudoephedrine Hydrochloride** 30 mg - Nasal Decongestant

**Uses** Temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies: ■ cough due to minor throat and bronchial irritation ■ nasal congestion ■ reduces swelling of nasal passages ■ runny nose ■ sneezing ■ itching of the nose or throat ■ itchy, watery eyes

**Warnings** Do not exceed recommended dosage.

### Do not use this product

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a physician or pharmacist before taking this product.

**Ask a physician before use if you have** ■ a cough that lasts or is chronic such as occurs with smoking, asthma or emphysema ■ a cough that occurs with too much phlegm (mucus) ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ trouble urinating due to an enlarged prostate gland ■ glaucoma ■ a breathing problem such as emphysema or chronic bronchitis. **Ask a physician or pharmacist before use if you are taking sedatives or tranquilizers.**

### When using this product

■ excitability may occur, especially in children ■ may cause marked drowsiness ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery ■ Stop use and ask a physician if ■ nervousness, dizziness, or sleeplessness occur ■ cough or nasal congestion persists for more than 1 week, tends to recur, or is accompanied by a fever, rash or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of the reach of children.**

In case of an accidental overdose, seek professional help or contact a Poison Control Center immediately.

**Directions** Do not exceed recommended dosage.

**Adults and children 12 years of age and over:**

2 teaspoonfuls (10 mL) every 6 hours,  
not to exceed 8 teaspoonfuls (40 mL) in 24 hours.

**Children 6 to under 12 years of age:**

1 teaspoonful (5 mL) every 6 hours, not to exceed  
4 teaspoonfuls (20 mL) in 24 hours.

**Children under 6 years of age:**

Consult a physician

### Other information

Store at 59° - 86° F (15° - 30° C)

**Inactive ingredients** Citric Acid, Glycerin, Grape Flavor, Propylene Glycol, Purified Water, Sodium Citrate, Sodium Saccharin, Sorbitol.

**Questions? Comments?** Call your physician for medical advice. Serious side effects associated with this product may be reported to this number. 1-888-206-5525  
8am - 5 pm M - F EST

Manufactured for:

**MAGNA**  
Pharmaceuticals, Inc.  
*Accountability.*

Louisville, KY 40299  
magnaweb.com  
888.206.5525