

“Hassle-Free Prescribing”

Z-TUSS™ AC 

**Cherry Flavor, Sugar Free, Alcohol Free
Antitussive • Antihistamine**

- **Hassle-free Prescribing or Recommending** (Schedule V)
- **Flexible Dosage:** Frequency and Volume Titration
- **Safety:** No Black Box Warning

Each Z-TUSS™ AC teaspoonful (5 ml) contains:

Codeine Phosphate 9mg
Chlorpheniramine Maleate 2mg

Directions

Do not exceed recommended dosage.

Adults and children over 12 years of age:	2 teaspoonsful (10 ml) every 4 to 6 hours, not to exceed 12 teaspoonfuls in 24 hours, or as directed by a physician
Children 6 to under 12 years of age:	1 teaspoonful (5 ml) every 4 to 6 hours, not to exceed 6 teaspoonfuls in 24 hours, or as directed by a physician
Children under 6 years of age	Consult a physician

Doses should not be repeated more than six times in a 24-hour period.

MAGNA
Pharmaceuticals, Inc.

Accountability •

www.magnaweb.com

NDC 58407-0920-04

NDC 58407-0920-04

Z-TUSS™ AC
Antitussive • Antihistamine

Drug Facts

Active ingredients

(in each 5mL teaspoonful)

Codeine Phosphate* 9 mg

Chlorpheniramine Maleate 2 mg

*(Warning: May be habit-forming)

Uses

Temporarily relieves runny nose, sneezing, itching of the nose or throat, and itchy, watery eyes due to hay fever or other upper respiratory allergies (allergic rhinitis). Temporarily relieves cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants and helps you cough less.

Warnings:

Do not exceed recommended dosage.

Do not take this product, unless directed by a doctor before use if you have

- a persistent or chronic cough such as occurs with smoking, asthma or emphysema
- a cough that occurs with too much phlegm (mucus)
- a chronic pulmonary disease, shortness of breath, or children who are taking other drugs
- a breathing problem such as emphysema or chronic bronchitis, or if you have glaucoma
- difficulty in urination due to enlargement of the prostate gland.

Ask a doctor before use if you are taking sedatives or tranquilizers

When using the product

- may cause marked drowsiness
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating
- excitability may occur, especially in children.

Stop use and ask a doctor if

- cough persists for more than 1 week, tends to recur, or is accompanied by a fever, rash or persistent headache; these could be signs of a serious condition.
- may cause or aggravate constipation.

If pregnant or breastfeeding, ask a health professional before use.

Keep this and other drugs out of the reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions:

Do not exceed recommended dosage.

Adults and children 12 years and over two teaspoonfuls (10 mL) every 4-6 hours

Children 6 to under 12 years of age one teaspoonful (5mL) every 4-6 hours

Children under 6 years of age Consult a doctor

A special measuring device should be used to give an accurate dose of this product to children under 6 years of age. Giving a higher dose than recommended by a doctor could result in serious effects for your child.

Other information

Store at 15°-30°C (59°-86°F)

Inactive ingredients

Cherry Flavor, Citric Acid, FD&C Red No. 40, Propylene Glycol, Purified Water, Saccharin Sodium, Sodium Benzoate, Sorbitol.

Questions or comments?

Call 1-888-206-5525

Manufactured for:
MAGNA Pharmaceuticals, Inc.
10801 Electron Drive, Suite 100
Louisville, KY 40299
www.magnaweb.com

